

THE ABUNDANCES OF ETHANE TO ACETYLENE IN THE ATMOSPHERES  
OF JUPITER AND SATURN

K. S. Noll and R. F. Knacke  
State University of New York at Stony Brook

A. T. Tokunaga  
University of Hawaii

J. H. Lacy  
University of Texas, Austin

S. Beck  
Northeastern University

E. Serabyn  
University of California, Berkeley

The presentation by Noll et al. is largely contained in a paper which appears in the special issue of *Icarus* (1986; 65, 257-263). The abstract of that paper is reproduced here.

Infrared spectra near  $780\text{ cm}^{-1}$  of Jupiter and Saturn have been obtained to determine the stratospheric abundances of ethane ( $\text{C}_2\text{H}_6$ ) and acetylene ( $\text{C}_2\text{H}_2$ ). Atmospheric models using Voyager thermal profiles and density profiles with constant mixing ratios result in the mixing ratios,  $X(\text{C}_2\text{H}_2) = 1.0(\pm 0.3) \times 10^{-7}$  and  $X(\text{C}_2\text{H}_6) = 5.5(\pm 1.5) \times 10^{-6}$  for Jupiter. The results for Saturn are  $X(\text{C}_2\text{H}_2) = 3.0(\pm 1.0) \times 10^{-7}$  and  $X(\text{C}_2\text{H}_6) = 7.0(\pm 1.5) \times 10^{-6}$ . The ratio of ethane to acetylene,  $n[\text{C}_2\text{H}_6]/n[\text{C}_2\text{H}_2]$ , is found to be insensitive to model atmosphere assumptions. The ratio is  $55 \pm 31$  for Jupiter and  $23 \pm 12$  for Saturn from models with uniform mixing ratios. Atmospheric models with density profiles adapted from theoretical photochemical models also result in a higher ratio of ethane to acetylene (by a factor of 2 at the 1 mbar level) on Jupiter. The lower abundance of acetylene on Jupiter suggests that the rate of vertical transport in the stratosphere may be more rapid on Saturn than on Jupiter.